

***Vamsapriya jinniuensis* sp. nov., and a first record of *Garnaudia elegans* from southern China**

YIN LING¹, HAO-HUA LI², JI-WEN XIA²,
XIU-GUO ZHANG^{2*}A, ZHUANG LI^{2*}B

¹ College of Life Sciences, Inner Mongolia University for Nationalities,
Tongliao, Inner Mongolia 028000, China

² Shandong Provincial Key Laboratory for Biology of Vegetable Diseases and Insect Pests,
College of Plant Protection, Shandong Agricultural University, Taian, 271018, China

*CORRESPONDENCE TO: ^A sdau613@163.com zhxg@sdau.edu.cn ^B liz552@126.com

ABSTRACT —A new species, *Vamsapriya jinniuensis*, is described and illustrated from specimens collected on dead stems in Hainan Province. The fungus is unique in producing catenate, phragmosporous, obclavate conidia on synnematos conidiophores with non-cicatrized, monotretic conidiogenous cells. *Garnaudia elegans* is newly recorded from China.

KEY WORDS —hyphomycetes, taxonomy, *Xylariaceae*

Introduction

During our ongoing survey of microfungi associated with woody debris in the forests of southern China, two specimens with the morphological characteristics of *Vamsapriya* Gawas & Bhat (Gawas & Bhat 2006) and *Garnaudia* Borowska (Borowska 1977) were collected on dead stems. *Vamsapriya jinniuensis* is described here as a new species, and *Garnaudia elegans* represents a new record for China. *Vamsapriya* currently includes six species: *V. bambusicola* D.Q. Dai & al., *V. breviconidiophora* Jun F. Li & al., *V. camagueyensis* (R.F. Castañeda) R.F. Castañeda & al., *V. indica* Gawas & Bhat, *V. khunkonensis* D.Q. Dai & al., and *V. mahabaleshwarensis* J. Pratibha & Bhat (Gawas & Bhat 2006, Pratibha & Bhat 2008, Dai & al. 2015, Castañeda-Ruiz & al. 2017, Hyde & al. 2017). Specimens are deposited in the

Herbarium of the Department of Plant Pathology, Shandong Agricultural University, Taian, China (HSAUP) and the Mycological Herbarium, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China (HMAS).

Vamsapriya jinniuensis Yin Ling & X.G. Zhang, sp. nov.

FIG. 1

MYCOBANK MB 826967

Differs from *Vamsapriya indica* by its longer, narrower, obclavate conidia.

TYPE: China, Hainan Province: Jinniuling, on dead stems of an unidentified broadleaf tree, 22 Feb. 2017, H.H. Li (Holotype, HSAUP H10242; isotype, HMAS 245647).

ETYMOLOGY: *jinniuensis*, in reference to the type locality, Jinniuling.

COLONIES on natural substrate effuse, dark brown to black. Mycelium mostly immersed in the substratum, composed of subhyaline, septate, branched, smooth hyphae. Synnemata erect, rigid, dark brown, composed of compact parallel conidiophores, ≤ 875 μm high, 36.5–103.5 μm diam. CONIDIOPHORES distinct, dark brown, smooth, septate, branched, 4–4.5 μm diam. CONIDIOGENOUS CELLS monotretic, uncitrized, integrated or discrete, terminal, clavate, slightly curved toward the exterior, $12\text{--}16 \times 3.8\text{--}4.5$ μm . CONIDIA dry, catenate, acrogenous, brown, smooth, simple, obclavate, 9–13-euseptate, constricted at the septa, $76\text{--}118.5 \times 2.5\text{--}5$ μm .

COMMENTS – *Vamsapriya* was established by Gawas & Bhat (2006), with *V. indica* as type species, to accommodate conidial fungi with catenate, phragmosporous conidia on synnematos conidiophores with non-citrized, monotretic, conidiogenous cells (Gawas & Bhat 2006). The genus is similar to *Didymobotryum* Sacc. and *Podosporium* Schwein. in having synnematos conidiomata and monotretic conidiogenous cells. However, conidia in *Didymobotryum* are catenate, ellipsoidal to cylindrical, and 1-septate, while in *Podosporium* they are solitary, obclavate, and multiseptate. *Vamsapriya* combines conidial morphology from both genera, bearing catenate, cylindrical to vermiform, multiseptate conidia.

Vamsapriya indica differs from *V. jinniuensis*, which has shorter, wider, cylindrical conidia with fewer septa ($10\text{--}80 \times 4\text{--}6$ μm , 2–12-euseptate; Gawas & Bhat 2006). The other five *Vamsapriya* species also all differ from *V. jinniuensis* by their shorter, wider conidia: *V. bambusicola* $8\text{--}45 \times 4.5\text{--}9.5$ μm (Dai & al. 2015); *V. breviconidiophora* $25.2\text{--}32.4 \times 7.9\text{--}10.2$ μm (Hyde & al. 2017); *V. camagueyensis* $45\text{--}65 \times 4\text{--}6$ μm (Castañeda-Ruiz & al. 2017); *V. khunkonensis* $17.5\text{--}35 \times 6\text{--}10$ μm (Dai & al. 2015); and *V. mahabaleshwarensis* $5\text{--}25 \times 4\text{--}9$ μm (Pratibha & Bhat 2008).




FIG. 1. *Vamsapriya jinniuensis* (holotype, HSAUP H10242). A. Synnema; B, C. Conidiophores and conidiogenous cells; D. Conidia.




FIG. 2. *Garraudia elegans* (HSAUP H10237). A. Conidiophore, conidiogenous cells, and conidia; B. Conidiogenous cells and conidia; C. Conidia; D. Conidiogenous cells.

Garnaudia elegans Borowska, Acta Mycol. 13(1): 170, 1977.

FIG. 2

CONIDIOPHORES macronematous, mononematous, brown, becoming paler towards the apex, erect, flexuous, simple or branched, up to 237.5 µm long, 2.5–4 µm wide towards the base. CONIDIOGENOUS CELLS monoblastic, discrete, determinate, ampuliform or lageniform, truncated, subhyaline or pale yellowish brown, solitary or in groups, 4.5–5.6 µm long, 2–3 µm wide at the apex of each stipe or branch. CONIDIA solitary, dry, acrogenous, simple, ellipsoidal, oval or piriform, 1-septate, basal cell 2× longer than apical cell, evenly pigmented, smooth, 6.5–8.5 × 3–5 µm, sometimes with a basal hyaline marginal frill.

SPECIMEN EXAMINED: CHINA, JIANGXI PROVINCE: Dayueshan, on dead stems of an unidentified broadleaf tree, 23 Dec. 2016, H.H. Li (HSAUP H10237).

COMMENTS – *Garnaudia* was established by Borowska (1977) with *G. elegans* as type species. It is mainly characterized by macronematous, simple or branched conidiophores, and monoblastic, discrete, determinate, terminal and intercalary, verticillate, ampuliform or lageniform conidiogenous cells that produce solitary, acrogenous, simple, brown, continuous or septate conidia.

We report *Garnaudia elegans* for the first time from China. Our Chinese specimen is similar to the Polish protologue description, but its conidiophores are slightly shorter than those of the type material (≤300 µm; Borowska 1977).

Acknowledgments

The authors express gratitude to Dr Li-Guo Ma and Dr Jian Ma for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by the National Natural Science Foundation of China (Nos. 31093440, 31230001, 31493010, 31493011, 31200013) and the Ministry of Science and Technology of the People's Republic of China (Nos. 2006FY120100).

Literature cited

- Borowska A. 1977. *Garnaudia elegans* gen. et sp. nov., and *Endophragmiella tenera* sp. nov., new dematiaceous hyphomycetes. Acta Mycologica 13(1): 169–174.
- Castañeda-Ruiz RE, Zhang XG, Li DW, Gusmão LFP, Pérez-Martínez S, Sosa D. 2017. Notes on *Vamsapriya* and *V. camagueyensis* comb. nov. Mycotaxon 132(3): 553–557. <https://doi.org/10.5248/132.553>
- Dai DQ, Bahkali AH, Li QR, Bhat DJ, Wijayawardene NN, Li WJ, Chuksatitrote E, Zhao RL, Xu JC, Hyde KD. 2015 [“2014”]. *Vamsapriya* (Xylariaceae) re-described, with two new species and molecular sequence data. Cryptogamie, Mycologie 35(4): 339–357. <https://doi.org/10.7872/crym.v35.iss4.2014.339>

- Gawas P, Bhat DJ. 2006 ["2005"]. *Vamsapriya indica* gen. et sp. nov., a bambusicolous, synnematosus fungus from India. *Mycotaxon* 94: 149–154.
- Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo A, Chethana KWT, & al. 2017. Fungal diversity notes 603–708: taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87: 1–235. <https://doi.org/10.1007/s13225-017-0391-3>
- Pratibha J, Bhat DJ. 2008. New and unusual hyphomycetes from Mahabaleshwar, India. *Mycotaxon* 105: 423–431.